

A Comparative Study of Variational Iteration and Adomian Decomposition Techniques for Solving Volterra Integro-Differential Equations

¹Kirtiwant P. Ghadle, ²Ahmed A. Hamoud, ³M.SH. Bani Issa

^{1,2}Department of Mathematics, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad, (M.S), India.

³P.E.T. Research Foundation Mandya, University of Mysore, Mysore-570 401 India.

Abstract—In this article, we present a comparative study between the Adomian decomposition and variational iteration methods. The study outlines the significant features of the two methods, for solving Volterra integro-differential equations. Moreover, we proved the existence and uniqueness results and convergence of the solution. Finally, some example are included to demonstrate the validity and applicability of the proposed techniques.

keywords: Adomian decomposition method, variational iteration method, Volterra integro-differential equation, approximate solution.

I. INTRODUCTION

In recent years, there has been a growing interest in the linear and nonlinear Volterra integro-differential equations which are a combination of differential and integral equations. The nonlinear Volterra integro-differential equations play an important role in many branches of nonlinear functional analysis and their applications in the theory of engineering, mechanics, physics, electrostatics, biology, chemistry and economics [1], [3], [5]. In this paper, we consider the Volterra integro-differential equations of the type:

$$\sum_{j=0}^k \xi_j(x) Z^{(j)}(x) = f(x) + \gamma \int_a^x K(x,t)G(Z(t))dt \quad (1)$$

with the initial conditions

$$Z^{(r)}(a) = b_r, \quad r = 0, 1, 2, \dots, (k-1), \quad (2)$$

where $Z^{(j)}(x)$ is the j^{th} derivative of the unknown function $Z(x)$ that will be determined, $K(x,t)$ is the kernel of the equation, $f(x)$ and $\xi_j(x)$ are an analytic function, G is nonlinear function of Z and a, b, γ , and b_r are real finite constants. Recently, many authors focus on the development of numerical and analytical techniques for integro-differential equations. For instance, we can remember the following works. Abbasbandy and Elyas [2] studied some applications on variational iteration method for solving system of nonlinear volterra integro-differential equations, Hamoud and Ghadle [6]

applied the hybrid methods for solving nonlinear Volterra-Fredholm integro-differential equations, Alao et al. [4] used Adomian decomposition and variational iteration methods for solving integro-differential equations, Yang and Hou [19] applied the Laplace decomposition method to solve the fractional integro-differential equations, Mittal and Nigam [14] applied the Adomian decomposition method to approximate solutions for fractional integro-differential equations, and Behzadi et al. [5] solved some class of nonlinear Volterra-Fredholm integro-differential equations by homotopy analysis method. Moreover, several authors have applied the Adomian decomposition method and the variational iteration method to find the approximate solutions of various types of integro-differential equations [6], [9], [10], [11], [14], [19].

The main objective of the present paper is to study the behavior of the solution that can be formally determined by semi-analytical approximated methods as the Adomian decomposition and variational iteration methods. Moreover, we proved the existence, uniqueness results and convergence of the solutions of the Volterra integro-differential equation (1).

II. DESCRIPTION OF THE METHODS

Some powerful methods have been focusing on the development of more advanced and efficient methods for integro-differential equations such as the Adomian decomposition method (ADM) [1], [3], [9], [11] and variational iteration method (VIM) [2], [4], [12], [13]. We will describe all these methods in this section:

A. Description of the ADM

Now, we can rewrite Eq.(1) in the form

$$\xi_k(x)Z^k(x) + \sum_{j=0}^{k-1} \xi_j(x)Z^j(x) = f(x) + \gamma \int_a^x K(x,t)G(Z(t))dt$$

Then

$$Z^k(x) = \frac{f(x)}{\xi_k(x)} + \gamma \int_a^x \frac{K(x,t)}{\xi_k(x)} G(Z(t))dt - \sum_{j=0}^{k-1} \frac{\xi_j(x)}{\xi_k(x)} Z^j(x).$$

To obtain the approximate solution, we integrating (k) -times in the interval $[a, x]$ with respect to x we obtain,

$$Z(x) = L^{-1} \left(\frac{f(x)}{\xi_k(x)} \right) + \sum_{r=0}^{k-1} \frac{1}{r!} (x-a)^r b_r + \gamma L^{-1} \left(\int_a^x \frac{K(x,t)}{\xi_k(x)} G(Z(t)) dt \right) - \sum_{j=0}^{k-1} L^{-1} \left(\frac{\xi_j(x)}{\xi_k(x)} Z_n^{(j)}(x) \right),$$

where L^{-1} is the multiple integration operator given as follows:

$$L^{-1}(\cdot) = \int_a^x \int_a^x \dots \int_a^x (\cdot) dx dx \dots dx \quad (k - \text{times}).$$

Now we apply ADM

$$G(Z(x)) = \sum_{n=0}^{\infty} A_n, \tag{3}$$

where $A_n; n \geq 0$ are the Adomian polynomials determined formally as follows:

$$A_n = \frac{1}{n!} \left[\frac{d^n}{d\mu^n} G \left(\sum_{i=0}^{\infty} \mu^i Z_i \right) \right]_{\mu=0}, \tag{4}$$

The Adomian polynomials were introduced in [15], [18], [19] as:

$$\begin{aligned} A_0 &= G(Z_0); \\ A_1 &= Z_1 G'(Z_0); \\ A_2 &= Z_2 G'(Z_0) + \frac{1}{2!} Z_1^2 G''(Z_0); \\ A_3 &= Z_3 G'(Z_0) + Z_1 Z_2 G''(Z_0) + \frac{1}{3!} Z_1^3 G'''(Z_0), \dots \end{aligned}$$

The standard decomposition technique represents the solution of Z as the following series:

$$Z = \sum_{i=0}^{\infty} Z_i. \tag{5}$$

By substituting (3) and (5) in Eq. (3) we have

$$\begin{aligned} \sum_{i=0}^{\infty} Z_i(x) &= L^{-1} \left(\frac{f(x)}{\xi_k(x)} \right) + \sum_{r=0}^{k-1} \frac{1}{r!} (x-a)^r b_r \\ &+ \gamma \sum_{i=0}^{\infty} L^{-1} \left(\int_a^x \frac{K(x,t)}{\xi_k(x)} A_i(t) dt \right) \\ &- \sum_{i=0}^{\infty} \sum_{j=0}^{k-1} L^{-1} \left(\frac{\xi_j(x)}{\xi_k(x)} Z_i^{(j)}(x) \right). \end{aligned} \tag{6}$$

The components Z_0, Z_1, Z_2, \dots are usually determined recursively by

$$\begin{aligned} Z_0 &= L^{-1} \left(\frac{f(x)}{\xi_k(x)} \right) + \sum_{r=0}^{k-1} \frac{1}{r!} (x-a)^r b_r, \\ Z_1 &= \gamma L^{-1} \left(\int_a^x \frac{K(x,t)}{\xi_k(x)} A_0(t) dt \right) - \sum_{j=0}^{k-1} L^{-1} \left(\frac{\xi_j(x)}{\xi_k(x)} Z_0^{(j)}(x) \right), \\ Z_{n+1} &= \gamma L^{-1} \left(\int_a^x \frac{K(x,t)}{\xi_k(x)} A_n(t) dt \right) - \sum_{j=0}^{k-1} L^{-1} \left(\frac{\xi_j(x)}{\xi_k(x)} Z_n^{(j)}(x) \right), \end{aligned}$$

Then, $Z(x) = \sum_{i=0}^n Z_i$ as the approximate solution.

B. Description of the VIM

This method is applied to solve a large class of linear and nonlinear problems with approximations converging rapidly to exact solutions. The main idea of this method is to construct a correction functional form using general Lagrange multipliers. These multipliers should be chosen such that its correction solution is superior to its initial approximation, called trial function. It is the best within the flexibility of trial functions. Accordingly, Lagrange multipliers can be identified by the variational theory [16], [17], [12]. A complete review of He's variational iteration method is available in [12], [13]. The initial approximation can be freely chosen with possible unknowns, which can be determined by imposing boundary/initial conditions. To illustrate, we consider the following general differential equation:

$$LZ(t) + NZ(t) = f(t), \tag{8}$$

where L is a linear operator, N is a nonlinear operator and $f(t)$ is inhomogeneous term. According to variational iteration method [4], the terms of a sequence Z_n are constructed such that this sequence converges to the exact solution. The terms Z_n are calculated by a correction functional as follows:

$$Z_{n+1}(t) = Z_n(t) + \int_0^t \lambda(\tau) (LZ_n(\tau) + NZ_n(\tau) - f(\tau)) d\tau. \tag{9}$$

The successive approximation $Z_n(t), n \geq 0$ of the solution $Z(t)$ will be readily obtained upon using the obtained Lagrange multiplier and by using any selective function Z_0 . The zeroth approximation Z_0 may be selected using any function that just satisfies at least the initial and boundary conditions. With λ determined, several approximations $Z_n(t), n \geq 0$ follow immediately.

The VIM has been shown to solve effectively, easily and accurately a large class of nonlinear problems with approximations converging rapidly to accurate solutions.

To obtain the approximation solution of IVB (1) – (2), according to the VIM, the iteration formula (9) can be written as follows:

$$\begin{aligned} Z_{n+1}(x) &= Z_n(x) + L^{-1} \left[\lambda(x) \left[\sum_{j=0}^k \xi_j(x) Z_n^{(j)}(x) - f(x) \right. \right. \\ &\quad \left. \left. - \gamma \int_a^x K(x,t) G(Z_n(t)) dt \right] \right], \end{aligned}$$

where L^{-1} is the multiple integration operator given as follows: Such that,

$$L^{-1}(\cdot) = \int_a^x \int_a^x \cdots \int_a^x (\cdot) dx dx \cdots dx \quad (k - \text{times}).$$

To find the optimal $\lambda(x)$, we proceed as follows:

$$\begin{aligned} \delta Z_{n+1}(x) &= \delta Z_n(x) + \delta L^{-1} \left[\lambda(x) \left[\sum_{j=0}^k \xi_j(x) Z_n^{(j)}(x) - f(x) \right. \right. \\ &\quad \left. \left. - \gamma \int_a^x K(x,t) G(Z_n(t)) dt \right] \right] \\ &= \delta Z_n(x) + \lambda(x) \delta Z_n(x) - L^{-1} \left[\delta Z_n(x) \lambda'(x) \right]. \end{aligned} \quad (10)$$

From Eq. (10), the stationary conditions can be obtained as follows:

$$\lambda'(x) = 0, \text{ and } 1 + \lambda(x)|_{x=t} = 0.$$

As a result, the Lagrange multipliers can be identified as $\lambda(x) = -1$ and by substituting in Eq. (10), the following iteration formula is obtained:

$$\begin{aligned} Z_0(x) &= L^{-1} \left[\frac{f(x)}{\xi_k(x)} \right] + \sum_{r=0}^{k-1} \frac{(x-a)^r}{r!} b_r, \\ Z_{n+1}(x) &= Z_n(x) - L^{-1} \left[\sum_{j=0}^k \xi_j(x) Z_n^{(j)}(x) - f(x) \right. \\ &\quad \left. - \gamma \int_a^x K(x,t) G(Z_n(t)) dt \right], n \geq 0. \end{aligned} \quad (11)$$

The term $\sum_{r=0}^{k-1} \frac{(x-a)^r}{r!} b_r$ is obtained from the initial conditions, $\xi_k(x) \neq 0$. Relation (11) will enable us to determine the components $Z_n(x)$ recursively for $n \geq 0$. Consequently, the approximation solution may be obtained by using

$$Z(t) = \lim_{n \rightarrow \infty} Z_n(t). \quad (12)$$

III. MAIN RESULTS

In this section, we shall give an existence and uniqueness results of Eq. (1), with the initial condition (2) and prove it [6], [7], [8].

We can write the equation (1) in the form of:

$$\begin{aligned} Z(x) &= L^{-1} \left[\frac{f(x)}{\xi_k(x)} \right] + \sum_{r=0}^{k-1} \frac{(x-a)^r}{r!} b_r \\ &\quad + \gamma L^{-1} \left[\int_a^x \frac{1}{\xi_k(x)} K(x,t) G(Z_n(t)) dt \right] \\ &\quad - L^{-1} \left[\sum_{j=0}^{k-1} \frac{\xi_j(x)}{\xi_k(x)} Z^{(j)}(x) \right]. \end{aligned} \quad (13)$$

$$\begin{aligned} L^{-1} \left[\int_a^x \frac{1}{\xi_k(x)} K(x,t) G(Z_n(t)) dt \right] &= \\ \int_a^x \frac{(x-t)^k}{k! \xi_k(x)} K(x,t) G(Z_n(t)) dt & \\ \sum_{j=0}^{k-1} L^{-1} \left[\frac{\xi_j(x)}{\xi_k(x)} \right] Z^{(j)}(x) &= \\ \sum_{j=0}^{k-1} \int_a^x \frac{(x-t)^{k-1} \xi_j(t)}{k-1! \xi_k(t)} Z^{(j)}(t) dt. & \end{aligned}$$

$$\Psi(x) = L^{-1} \left[\frac{f(x)}{\xi_k(x)} \right] + \sum_{r=0}^{k-1} \frac{(x-a)^r}{r!} b_r.$$

Before starting and proving the main results, we introduce the following hypotheses:

- (H1) There exist two constants α and $\gamma_j > 0, j = 0, 1, \dots, k$ such that, for any $Z_1, Z_2 \in C(J, \mathbb{R})$

$$|G(Z_1) - G(Z_2)| \leq \alpha |Z_1 - Z_2|$$

and

$$|D^j(Z_1) - D^j(Z_2)| \leq \gamma_j |Z_1 - Z_2|,$$

we suppose that the nonlinear terms $G(Z(x))$ and $D^j(Z) = \left(\frac{d^j}{dx^j}\right)Z(x) = \sum_{i=0}^{\infty} \gamma_{ij}$, (D^j is a derivative operator), $j = 0, 1, \dots, k$, are Lipschitz continuous.

- (H2) we suppose that for all $a \leq t \leq x \leq b$, and $j = 0, 1, \dots, k$:

$$\left| \frac{\gamma(x-t)^k K(x,t)}{k! \xi_k(x)} \right| \leq \theta_1, \quad \left| \frac{\gamma(x-t)^k K(x,t)}{k!} \right| \leq \theta_2,$$

$$\left| \frac{(x-t)^{k-1} \xi_j(t)}{(k-1)! \xi_k(t)} \right| \leq \theta_3, \quad \left| \frac{(x-t)^{k-1} \xi_j(t)}{(k-1)!} \right| \leq \theta_4.$$

- (H3) There exist three functions θ_3^*, θ_4^* , and $\gamma^* \in C(D, \mathbb{R}^+)$, the set of all positive function continuous on $D = \{(x,t) \in \mathbb{R} \times \mathbb{R} : 0 \leq t \leq x \leq 1\}$ such that:

$$\theta_3^* = \max |\theta_3|, \quad \theta_4^* = \max |\theta_4|, \quad \text{and } \gamma^* = \max |\gamma_j|.$$

- (H4) $\Psi(x)$ is bounded function for all x in $J = [a, b]$.

Theorem 1. Assume that (H1)–(H4) hold. If

$$0 < \psi = (\alpha \theta_1 + k \gamma^* \theta_3^*)(b-a) < 1, \quad (14)$$

Then there exists a unique solution $Z(x) \in C(J)$ to IVB (1)–(2).

Proof. Let Z_1 and Z_2 be two different solutions of IVB (1) – (2)., then

$$\begin{aligned} |Z_1 - Z_2| &= \left| \int_a^x \frac{\gamma(x-t)^k K(x,t)}{\xi_k(x)k!} [G(Z_1) - G(Z_2)] dt \right. \\ &\quad \left. - \sum_{j=0}^{k-1} \int_a^x \frac{(x-t)^{k-1} \xi_j(t)}{\xi_k(t)(k-1)!} [D^j(Z_1) - D^j(Z_2)] dt \right| \\ &\leq \int_a^x \left| \frac{\gamma(x-t)^k K(x,t)}{\xi_k(x)k!} \right| |G(Z_1) - G(Z_2)| dt \\ &\quad - \sum_{j=0}^{k-1} \int_a^x \left| \frac{(x-t)^{k-1} \xi_j(t)}{\xi_k(t)(k-1)!} \right| |D^j(Z_1) - D^j(Z_2)| dt \\ &\leq (\alpha\theta_1 + k\gamma^*\theta_3^*)(b-a)|Z_1 - Z_2|, \end{aligned}$$

we get $(1-\psi)|Z_1 - Z_2| \leq 0$. Since $0 < \psi < 1$, so $|Z_1 - Z_2| = 0$. therefore, $Z_1 = Z_2$ and the proof is completed. \square

Theorem 2. Suppose that (H1)–(H4), and If $0 < \psi < 1$, hold, the series solution $Z(x) = \sum_{m=0}^{\infty} Z_m(x)$ and $\|Z_1\|_{\infty} < \infty$ obtained by the m -order deformation is convergent, then it converges to the exact solution of the Volterra integro-differential equation (1) – (2).

Proof. Denote as $(C[0,1], \|\cdot\|)$ the Banach space of all continuous functions on J , with $|Z_1(x)| \leq \infty$ for all x in J .

Frist we define the sequence of partial sums v_n , let v_n and v_m be arbitrary partial sums with $n \geq m$. We are going to prove that $v_n = \sum_{i=0}^n Z_i(x)$ is a Cauchy sequence in this Banach space:

$$\begin{aligned} \|v_n - v_m\|_{\infty} &= \max_{\forall x \in J} |v_n - v_m| \\ &= \max_{\forall x \in J} \left| \sum_{i=0}^n Z_i(x) - \sum_{i=0}^m Z_i(x) \right| \\ &= \max_{\forall x \in J} \left| \sum_{i=m+1}^n Z_i(x) \right| \\ &= \max_{\forall x \in J} \left| \sum_{i=m+1}^n \int_a^x \frac{\gamma(x-t)^k K(x,t)}{k! \xi_k(x)} A_{i-1} dt \right. \\ &\quad \left. - \sum_{j=0}^{k-1} \int_a^x \frac{\xi_j(t)(x-t)^{k-1}}{(k-1)! \xi_k(t)} \gamma_{(i-1)_J} dt \right| \\ &= \max_{\forall x \in J} \left| \int_a^x \frac{\gamma(x-t)^k K(x,t)}{k! \xi_k(x)} \left(\sum_{i=m}^{n-1} A_i \right) dt \right. \\ &\quad \left. - \sum_{j=0}^{k-1} \int_a^x \frac{\xi_j(t)(x-t)^{k-1}}{(k-1)! \xi_k(t)} \left(\sum_{i=m}^{n-1} \gamma_{i_j} dt \right) \right|. \end{aligned}$$

From (4), we have

$$\begin{aligned} \sum_{i=m}^{n-1} A_i &= G(v_{n-1}) - G(v_{m-1}), \\ \sum_{i=m}^{n-1} \gamma_i &= D^j(v_{n-1}) - D^j(v_{m-1}). \end{aligned}$$

$$\begin{aligned} \|v_n - v_m\|_{\infty} &= \max_{\forall x \in J} \left| \int_a^x \frac{\gamma(x-t)^k K(x,t)}{k! \xi_k(x)} \right. \\ &\quad \times [G(v_{n-1}) - G(v_{m-1})] dt \\ &\quad \left. - \sum_{j=0}^{k-1} \int_a^x \frac{\xi_j(t)(x-t)^{k-1}}{(k-1)! \xi_k(t)} [D^j(v_{n-1}) - D^j(v_{m-1})] dt \right| \\ &\leq \max_{\forall x \in J} \int_a^x \left| \frac{\gamma(x-t)^k K(x,t)}{k! \xi_k(x)} \right| |G(v_{n-1}) - G(v_{m-1})| dt \\ &\quad + \sum_{j=0}^{k-1} \int_a^x \left| \frac{\xi_j(t)(x-t)^{k-1}}{(k-1)! \xi_k(t)} \right| |D^j(v_{n-1}) - D^j(v_{m-1})| dt \end{aligned}$$

Let $n = m + 1$, then

$$\begin{aligned} \|v_n - v_m\|_{\infty} &\leq \psi \|v_m - v_{m-1}\|_{\infty} \leq \psi^2 \|v_{m-1} - v_{m-2}\|_{\infty} \\ &\leq \dots \leq \psi^m \|v_1 - v_0\|_{\infty}, \end{aligned}$$

so,

$$\begin{aligned} \|v_n - v_m\|_{\infty} &\leq \|v_{m+1} - v_m\|_{\infty} + \|v_{m+2} - v_{m+1}\|_{\infty} \\ &\quad + \dots + \|v_n - v_{n-1}\|_{\infty} \\ &\leq [\psi^m + \psi^{m+1} + \dots + \psi^{n-1}] \|v_1 - v_0\|_{\infty} \\ &\leq \psi^m [1 + \psi + \psi^2 + \dots + \psi^{n-m-1}] \|v_1 - v_0\|_{\infty} \\ &\leq \psi^m \left(\frac{1 - \psi^{n-m}}{1 - \psi} \right) \|Z_1\|_{\infty}. \end{aligned}$$

Since $0 < \psi < 1$, we have $(1 - \psi^{n-m}) < 1$, then

$$\|v_n - v_m\|_{\infty} \leq \frac{\psi^m}{1 - \psi} \|Z_1\|_{\infty}.$$

But $|Z_1(x)| < \infty$, so, as $m \rightarrow \infty$, then $\|v_n - v_m\|_{\infty} \rightarrow 0$.

We conclude that v_n is a Cauchy sequence in $C[0,1]$, therefore $Z = \lim_{n \rightarrow \infty} Z_n$.

Then, the series is convergence and the proof is complete. \square

Theorem 3. If problem (1)–(2) has a unique solution, then the solution $Z_n(x)$ obtained from the recursive relation (11) using VIM converges when $0 < \phi = (\alpha\theta_2 + k\gamma^*\theta_4^*)(b-a) < 1$.

Proof. We have from equation (11):

$$\begin{aligned} Z_{n+1}(x) - Z(x) &= Z_n(x) - Z(x) \\ &\quad - \left(L^{-1} \left[\sum_{j=0}^k \xi_j(x) [Z_n^{(j)}(x) - Z^{(j)}(x)] \right] \right. \\ &\quad \left. - L^{-1} \left[\gamma \int_a^x K(x,t) [G(Z_n(t)) - G(Z(t))] dt \right] \right). \end{aligned}$$

If we set, $\xi_k(x) = 1$, and $W_{n+1}(x) = Z_{n+1}(x) - Z(x)$, $W_n(x) = Z_n(x) - Z(x)$ since $W_n(a) = 0$, then

$$\begin{aligned} W_{n+1}(x) &= W_n(x) + \int_a^x \frac{\gamma K(x,t)(x-t)^k}{k!} [G(Z_n(t)) - G(Z(t))] dt \\ &\quad - \sum_{j=0}^{k-1} \int_a^x \frac{\lambda_1 \xi_j(t)(x-t)^{k-1}}{(k-1)!} [D^j(Z_n(t)) - D^j(Z(t))] dt \\ &\quad - (W_n(x) - W_n(a)). \end{aligned}$$

Therefore,

$$\begin{aligned}
 |W_{n+1}(x)| &\leq \int_a^x \left| \frac{\gamma K(x,t)(x-t)^k}{k!} \right| |W_n| \alpha dt \\
 &+ \sum_{j=0}^{k-1} \int_a^x \left| \frac{\gamma \xi_j(t)(x-t)^{k-1}}{(k-1)!} \right| \max |\gamma_j| |W_n| dt \\
 &\leq |W_n| \left[\int_a^x \alpha \theta_5 dt + \sum_{j=0}^{k-1} \int_a^x \theta_4^* | \max |\gamma_j| \right] \\
 &\leq |W_n| (\alpha \theta_2 + k \gamma^* \theta_4^*) (b-a) = |W_n| \phi.
 \end{aligned}$$

Hence,

$$\|W_{n+1}\| = \max_{\forall x \in J} |W_{n+1}(x)| \leq \phi \max_{\forall x \in J} |W_n(x)| = \phi \|W_n\|.$$

Since $0 < \phi < 1$, then $\|W_n\| \rightarrow 0$. So, the series converges and the proof is complete. \square

IV. NUMERICAL RESULTS

In this section, we present the semi-analytical techniques based on ADM, and VIM to solve Volterra integro-differential equations. To show the efficiency of the present methods for our problem in comparison with the exact solution we report absolute error.

Example 1. Consider the following Volterra integro-differential equation.

$$Z''(x) = xe^x + 1 - \int_0^x e^{x-s} Z(s) ds,$$

with the initial condition

$$Z(0) = 0, Z'(0) = 1,$$

and the the exact solution is $Z(x) = e^x - 1$.

TABLE I
NUMERICAL RESULTS OF THE EXAMPLE 1.

x	Exact solution	ADM	VIM
0.1	0.105170918	0.1051709278	0.1051709640
0.2	0.221402758	0.2214027799	0.2214028205
0.3	0.349858808	0.3498587427	0.3498586872
0.4	0.491824698	0.4918247080	0.4918245703
0.5	0.648721271	0.6487212399	0.6487213115
0.6	0.822118800	0.8221188288	0.8221188591
0.7	1.013752707	1.0137527321	1.0137527691
0.8	1.225540928	1.2255410371	1.2255409831
0.9	1.459603111	1.4596032701	1.4596031481
1.0	1.718281828	1.7182822951	1.7182819261

Example 2. Consider the following Volterra integro-differential equation.

$$Z'(x) = -1 + \int_0^x Z^2(s) ds,$$


with the initial condition

$$Z(0) = 0,$$

and the the exact solution is $Z(x) = -x$.

TABLE II
NUMERICAL RESULTS OF THE EXAMPLE 2.

x	Exact solution	ADM	VIM
0.1	-0.1000000000	-0.09999166707	-0.09999166707
0.2	-0.2000000000	-0.19986671750	-0.19986671750
0.3	-0.3000000000	-0.29932586690	-0.29932586690
0.4	-0.4000000000	-0.39787315100	-0.39787315100
0.5	-0.5000000000	-0.49482250800	-0.49482250780
0.6	-0.6000000000	-0.58931009410	-0.58931009270
0.7	-0.7000000000	-0.68031385970	-0.68031384920
0.8	-0.8000000000	-0.76668145931	-0.76668140030
0.9	-0.9000000000	-0.84716692600	-0.84716665890
1.0	-1.0000000000	-0.92047571081	-0.92047468820



V. COMPARISON AMONG THE METHODS

The comparison among of the methods, it can be seen from the results of the above example:

- The methods are powerful, efficient and give approximations of higher accuracy. Also, they can produce closed-form solutions if they exist.
- Although the results obtained by these methods when applied to Volterra integro-differential equations are the same approximately. VIM is seen to be much easier and more convenient than the ADM.
- One advantage of VIM is that the initial solution can be freely chosen with some unknown parameters. An interesting point about this method is that with few number of iterations, or even in some cases with only one iteration, it can produce a very accurate approximate solution.
- The VIM has a more rapid convergence than the ADM. Also, the number of computations in VIM is less than the ones in ADM.

VI. CONCLUSIONS

We present a comparative study between the Adomian decomposition and variational iteration methods for solving Volterra integro-differential equations. From the computational viewpoint, the variational iteration method is more efficient, convenient and easy to use. Moreover, we proved the existence and uniqueness results and convergence of the solution. The methods are very powerful and efficient in finding analytical as well as numerical solutions for wide classes of linear and non-linear Volterra integro-differential equations. The convergence theorems and the numerical results establish the precision and efficiency of the proposed techniques.

REFERENCES

- [1] Abbaoui, K. and Cherruault, Y. Convergence of Adomian's method applied to nonlinear equations, *Math. Comput. Modelling*, **20**(9) (1994), 69–73.
- [2] Abbasbandy, S. and Elyas, S. Application of variational iteration method for system of nonlinear volterra integro-differential equations, *Mathematics and computational applications*, **2**(14) (2009), 147–158.
- [3] Adomian, G. A review of the decomposition method in applied mathematics, *J. Math. Anal. Appl.*, **135**(2) (1988), 501–544.
- [4] Alao, S. Akinboro I, F. Akinpelu, F. and Oderinu, R. Numerical solution of integro-differential equation using Adomian decomposition and variational iteration methods, *IOSR Journal of Mathematics*, **10**(4) (2014), 18–22.
- [5] Behzadi, S. Abbasbandy, S. Allahviranloo, T. and Yildirim, A. Application of homotopy analysis method for solving a class of nonlinear Volterra-Fredholm integro-differential equations, *J. Appl. Anal. Comput.* **2**(2) (2012), 127–136.
- [6] Hamoud, A.A. and Ghadle, K.P. The approximate solutions of fractional Volterra-Fredholm integro-differential equations by using analytical techniques, *Probl. Anal. Issues Anal.*, **7**(25) (2018), 41–58.
- [7] Burton, T.A. Integro-differential equations, compact maps, positive kernels, and Schaefer's fixed point theorem, *Nonlinear Dyn. Syst. Theory*, **17**(1) (2017), 19–28.
- [8] Burton, T.A. Existence and uniqueness results by progressive contractions for integro-differential equations, *Nonlinear Dyn. Syst. Theory*, **16**(4) (2016), 366–371.
- [9] Hamoud, A.A. and Ghadle, K.P. The reliable modified of Laplace Adomian decomposition method to solve nonlinear interval Volterra-Fredholm integral equations, *Korean J. Math.* **25**(3) (2017), 323–334.
- [10] Hamoud, A.A. and Ghadle, K.P. Existence and uniqueness theorems for fractional Volterra-Fredholm integro-differential equations, *Int. J. Appl. Math.* **31**(3) (2018), 333–348.
- [11] Hamoud, A.A. and Ghadle, K.P. Modified Adomian decomposition method for solving fuzzy Volterra-Fredholm integral equations, *J. Indian Math. Soc.* **85**(1-2) (2018), 52–69.
- [12] He, J.H. A variational approach to nonlinear problems and its application, *Mech. Applic.* **20**(1) (1998), 30–34.
- [13] He, J.H. and Wang, S.Q. Variational iteration method for solving integro-differential equations, *Phys. Lett. A* **367** (2007), 188–191.
- [14] Mittal, R. and Nigam, R. Solution of fractional integro-differential equations by Adomian decomposition method, *Int. J. Appl. Math. Mech.*, **4**, 2 (2008), 87–94.
- [15] Wazwaz, A.M. A new algorithm for calculating Adomian polynomials for nonlinear operators, *Appl. Math. and Comput.* **111** (2000), 53–69.
- [16] Wazwaz, A.M. A comparison between variational iteration method and Adomian decomposition method, *Journal of computational and applied Mathematics*, **207** (2007), 129–136.
- [17] Wazwaz, A.M. The variational iteration method for solving linear and non-linear Volterra integral and integro-differential equations, *Int. J. Comput. Math.* **87**(5) (2010), 1131–1141.
- [18] Wazwaz, A.M. *Linear and Nonlinear Integral Equations Methods and Applications*, Springer Heidelberg Dordrecht London New York, 2011.
- [19] Yang, C. and Hou, J. Numerical solution of integro-differential equations of fractional order by Laplace decomposition method, *Wseas Trans. Math.*, **12**(12) (2013), 1173–1183.