

# $\alpha$ Generalized Closed Sets in Neutrosophic Topological Spaces

D. Jayanthi

Department of Mathematics

Avinashilingam (Deemed to be) University, Tamilnadu, India

**Abstract:** In this paper a new concept of neutrosophic closed sets called neutrosophic  $\alpha$  generalized closed sets is introduced and their properties are thoroughly studied and analyzed. Some new interesting theorems based on the newly introduced set are presented.

**Mathematics Subject Classification (2000):** 54A40, 03E72

**Keywords:** Neutrosophic sets, neutrosophic topology, neutrosophic  $\alpha$  generalized closed sets.

## 1. Introduction

The concept of neutrosophic sets was first introduced by Floretin Smarandache [3] in 1999 which is a generalization of intuitionistic fuzzy sets by Atanassov [1]. A. A. Salama and S. A. Alblowi [6] introduced the concept of neutrosophic topological spaces after Coker [2] introduced intuitionistic fuzzy topological spaces. Further the basic sets like semi open sets, pre open sets,  $\alpha$  open sets and semi- $\alpha$  open sets are introduced in neutrosophic topological spaces and their properties are studied by various authors [4,5]. The purpose of this paper is to introduce and analyze a new concept of neutrosophic closed sets called neutrosophic  $\alpha$  generalized closed sets.

## 2. Preliminaries:

Here in this paper the neutrosophic topological space is denoted by  $(X, \tau)$ . Also the neutrosophic interior, neutrosophic closure of a neutrosophic set  $A$  are denoted by  $NInt(A)$  and  $NCl(A)$ . The complement of a neutrosophic set  $A$  is denoted by  $C(A)$  and the empty and whole sets are denoted by  $0_N$  and  $1_N$  respectively.

**Definition 2.1:** Let  $X$  be a non-empty fixed set. A neutrosophic set (NS)  $A$  is an object having the form  $A = \{\langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle : x \in X\}$  where  $\mu_A(x)$ ,  $\sigma_A(x)$  and  $\nu_A(x)$  represent the degree of membership, degree of indeterminacy and the degree of non-

membership respectively of each element  $x \in X$  to the set  $A$ .

A Neutrosophic set  $A = \{\langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle : x \in X\}$  can be identified as an ordered triple  $\langle \mu_A, \sigma_A, \nu_A \rangle$  in  $]0, 1^+[$  on  $X$ .

**Definition 2.2:** Let  $A = \langle \mu_A, \sigma_A, \nu_A \rangle$  be a NS on  $X$ , then the complement  $C(A)$  may be defined as

1.  $C(A) = \{\langle x, 1 - \mu_A(x), 1 - \nu_A(x) \rangle : x \in X\}$
2.  $C(A) = \{\langle x, \nu_A(x), \sigma_A(x), \mu_A(x) \rangle : x \in X\}$
3.  $C(A) = \{\langle x, \nu_A(x), 1 - \sigma_A(x), \mu_A(x) \rangle : x \in X\}$

Note that for any two neutrosophic sets  $A$  and  $B$ ,

4.  $C(A \cup B) = C(A) \cap C(B)$
5.  $C(A \cap B) = C(A) \cup C(B)$

**Definition 2.3:** For any two neutrosophic sets  $A = \{\langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle : x \in X\}$  and  $B = \{\langle x, \mu_B(x), \sigma_B(x), \nu_B(x) \rangle : x \in X\}$  we may have

1.  $A \subseteq B \Leftrightarrow \mu_A(x) \leq \mu_B(x), \sigma_A(x) \leq \sigma_B(x)$  and  $\nu_A(x) \geq \nu_B(x) \forall x \in X$
2.  $A \subseteq B \Leftrightarrow \mu_A(x) \leq \mu_B(x), \sigma_A(x) \geq \sigma_B(x)$  and  $\nu_A(x) \geq \nu_B(x) \forall x \in X$
3.  $A \cap B = \langle x, \mu_A(x) \wedge \mu_B(x), \sigma_A(x) \wedge \sigma_B(x), \nu_A(x) \vee \nu_B(x) \rangle$
4.  $A \cap B = \langle x, \mu_A(x) \wedge \mu_B(x), \sigma_A(x) \vee \sigma_B(x), \nu_A(x) \vee \nu_B(x) \rangle$
5.  $A \cup B = \langle x, \mu_A(x) \vee \mu_B(x), \sigma_A(x) \vee \sigma_B(x), \nu_A(x) \wedge \nu_B(x) \rangle$
6.  $A \cup B = \langle x, \mu_A(x) \vee \mu_B(x), \sigma_A(x) \wedge \sigma_B(x), \nu_A(x) \wedge \nu_B(x) \rangle$

**Definition 2.4:** A neutrosophic topology (NT) on a non-empty set  $X$  is a family  $\tau$  of neutrosophic subsets in  $X$  satisfies the following axioms:

- (NT<sub>1</sub>)  $0_N, 1_N \in \tau$
- (NT<sub>2</sub>)  $G_1 \cap G_2 \in \tau$  for any  $G_1, G_2 \in \tau$
- (NT<sub>3</sub>)  $\cup G_i \in \tau \forall \{G_i : i \in J\} \subseteq \tau$

In this case the pair  $(X, \tau)$  is a neutrosophic topological space (NTS) and any neutrosophic set in  $\tau$  is known as a neutrosophic open set (NOS) in  $X$ . A neutrosophic set  $A$  is a neutrosophic closed set (NCS)

if and only if its complement  $C(A)$  is a neutrosophic open set in  $X$ .

Here the empty set  $(0_N)$  and the whole set  $(1_N)$  may be defined as follows:

- (0<sub>1</sub>)  $0_N = \{\langle x, 0, 0, 1 \rangle : x \in X\}$
- (0<sub>2</sub>)  $0_N = \{\langle x, 0, 1, 1 \rangle : x \in X\}$
- (0<sub>3</sub>)  $0_N = \{\langle x, 0, 1, 0 \rangle : x \in X\}$
- (0<sub>4</sub>)  $0_N = \{\langle x, 0, 0, 0 \rangle : x \in X\}$
- (1<sub>1</sub>)  $1_N = \{\langle x, 1, 0, 0 \rangle : x \in X\}$
- (1<sub>2</sub>)  $1_N = \{\langle x, 1, 0, 1 \rangle : x \in X\}$
- (1<sub>3</sub>)  $1_N = \{\langle x, 1, 1, 0 \rangle : x \in X\}$
- (1<sub>4</sub>)  $1_N = \{\langle x, 1, 1, 1 \rangle : x \in X\}$

**Definition 2.5:** Let  $(X, \tau)$  be a NTS and  $A = \{\langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle : x \in X\}$  be a NS in  $X$ . Then the neutrosophic interior and the neutrosophic closure of  $A$  are defined by

$$NInt(A) = \cup \{G : G \text{ is an NOS in } X \text{ and } G \subseteq A\}$$

$$NCl(A) = \cap \{K : K \text{ is an NCS in } X \text{ and } A \subseteq K\}$$

Note that for any NS  $A$ ,  $NCl(C(A)) = C(NInt(A))$  and  $NInt(C(A)) = C(NCl(A))$ .

**Definition 2.6:** A NS  $A$  of a NTS  $X$  is said to be

- (i) a neutrosophic pre-open set (NP-OS) if  $A \subseteq NInt(NCl(A))$
- (ii) a neutrosophic semi-open set (NS-OS) if  $A \subseteq NCl(NInt(A))$
- (iii) a neutrosophic  $\alpha$ -open set ( $N\alpha$ -OS) if  $A \subseteq NInt(NCl(NInt(A)))$
- (iv) a neutrosophic semi- $\alpha$ -open set ( $NS_{\alpha}$ -OS) if  $A \subseteq NCl(\alpha NInt(A))$

**Definition 2.7:** A NS  $A$  of a NTS  $X$  is said to be

- (i) A neutrosophic pre-closed set (NP-CS) if  $NCl(NInt(A)) \subseteq A$
- (ii) A neutrosophic semi-closed set (NS-CS) if  $NInt(NCl(A)) \subseteq A$
- (iii) A neutrosophic  $\alpha$ -closed set ( $N\alpha$ -CS) if  $NCl(NInt(NCl(A))) \subseteq A$
- (iv) A neutrosophic semi- $\alpha$ -closed set ( $NS_{\alpha}$ -CS) if  $NInt(\alpha NCl(A)) \subseteq A$

### 3. $\alpha$ generalized closed sets in neutrosophic topological spaces

In this section we introduce neutrosophic  $\alpha$  closure, neutrosophic  $\alpha$  interior and  $\alpha$  generalized closed set and its respective open set in neutrosophic topological spaces and discuss some of their properties.

**Definition 3.1:** A NS  $A$  in a NTS  $X$  is said to be a neutrosophic regular closed set (NRCS) if

$NCl(NInt(A)) = A$  and neutrosophic regular open set if  $NInt(NCl(A)) = A$ .

**Definition 3.2:** A NS  $A$  in a NTS  $X$  is said to be a neutrosophic  $\beta$  closed set ( $N\beta$ CS) if  $NInt(NCl(NInt(A))) \sqcap A$  and neutrosophic  $\beta$  open set if  $A \sqcap NCl(NInt(NCl(A)))$

**Definition 3.3:** Let  $A$  be a NS of a NTS  $(X, \tau)$ . Then the neutrosophic  $\alpha$  interior and the neutrosophic  $\alpha$  closure are defined as

$$N_{\alpha}Int(A) = \cup \{G : G \text{ is a } N\alpha\text{-OS in } X \text{ and } G \subseteq A\}$$

$$N_{\alpha}Cl(A) = \cap \{K : K \text{ is a } N\alpha\text{-CS in } X \text{ and } A \subseteq K\}$$

**Result 3.4:** Let  $A$  be a NS in  $X$ . Then  $N_{\alpha}Cl(A) = A \cup NCl(NInt(NCl(A)))$ .

**Proof:** Since  $N_{\alpha}Cl(A)$  is a  $N\alpha$ -CS,  $NCl(NInt(NCl(N_{\alpha}Cl(A)))) \subseteq N_{\alpha}Cl(A)$  and  $A \cup NCl(NInt(NCl(A))) \subseteq A \cup NCl(NInt(NCl(N_{\alpha}Cl(A)))) \subseteq A \cup N_{\alpha}Cl(A) = N_{\alpha}Cl(A)$  -----(i). Now  $NCl(NInt(NCl(A \cup NCl(NInt(NCl(A))))) \subseteq NCl(NInt(NCl(A \cup NCl(A)))) = NCl(NInt(NCl(NCl(A)))) = NCl(NInt(NCl(A))) \subseteq A \cup NCl(NInt(NCl(A)))$ . Therefore  $A \cup NCl(NInt(NCl(A)))$  is a  $N\alpha$ -CS in  $X$  and hence  $N_{\alpha}Cl(A) \subseteq A \cup NCl(NInt(NCl(A)))$  -----(ii). From (i) and (ii),  $N_{\alpha}Cl(A) = A \cup NCl(NInt(NCl(A)))$ .

**Definition 3.5:** A NS  $A$  in a NTS  $X$  is said to be a neutrosophic  $\alpha$  generalized closed set ( $N_{\alpha g}$ CS) if  $N_{\alpha}Cl(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is a NOS in  $X$ . The complement  $C(A)$  of a  $N_{\alpha g}$ CS  $A$  is a  $N_{\alpha g}$ OS in  $X$ .

**Example 3.6:** Let  $X = \{a, b\}$  and  $\tau = \{0_N, A, B, 1_N\}$  where  $A = \langle x, (0.5, 0.6), (0.3, 0.2), (0.4, 0.1) \rangle$  and  $B = \langle x, (0.4, 0.4), (0.4, 0.3), (0.5, 0.4) \rangle$ . Then  $\tau$  is a NT. Here  $\mu_A(a) = 0.5$ ,  $\mu_A(b) = 0.6$ ,  $\sigma_A(a) = 0.3$ ,  $\sigma_A(b) = 0.2$ ,  $\nu_A(a) = 0.4$  and  $\nu_A(b) = 0.1$ . Also  $\mu_B(a) = 0.4$ ,  $\mu_B(b) = 0.4$ ,  $\sigma_B(a) = 0.4$ ,  $\sigma_B(b) = 0.3$ ,  $\nu_B(a) = 0.5$  and  $\nu_B(b) = 0.4$ . Let  $M = \langle x, (0.5, 0.4), ((0.4, 0.4), (0.4, 0.5)) \rangle$  be any NS in  $X$ . Then  $M \subseteq A$  where  $A$  is a NOS in  $X$ . Now  $N_{\alpha}Cl(M) = M \cup C(B) = C(B) \subseteq A$ . Therefore  $M$  is a  $N_{\alpha g}$ -CS in  $X$ .

**Proposition 3.7:** Every NCS  $A$  is a  $N_{\alpha g}$ -CS in  $X$  but not conversely in general.

**Proof:** Let  $A \subseteq U$  where  $U$  is a NOS in  $X$ . Now  $N_{\alpha}Cl(A) = A \cup NCl(NInt(NCl(A))) \subseteq A \cup NCl(A) = A \cup A = A \subseteq U$ , by hypothesis. Therefore  $A$  is a  $N_{\alpha g}$ -CS in  $X$ .

**Example 3.8:** In Example 3.6,  $M$  is a  $N_{\alpha g}$ -CS in  $X$  but not a NCS in  $X$  as  $NCl(M) = C(B) \neq M$ .

**Remark 3.9:** Every NS-CS and every  $N_{\alpha g}$ -CS in a NTS  $X$  are independent to each other in general.

**Example 3.10:** In Example 3.6,  $M$  is a  $N_{\alpha g}$ -CS but not a NS-CS as  $NInt(NCl(M)) = B \not\subseteq M$ .

**Example 3.11:** Let  $X = \{a, b\}$  and  $\tau = \{0_N, A, B, C, 1_N\}$ , where  $A = \langle x, (0.5, 0.4), (0.3, 0.2), (0.5, 0.6) \rangle$ ,  $B = \langle x, (0.8, 0.7), (0.4, 0.3), (0.2, 0.3) \rangle$  and  $C = \langle x, (0.2, 0.1), (0.3, 0.2), (0.8, 0.9) \rangle$ . Then  $\tau$  is a NT. Let  $M = \langle x, (0.5, 0.3), (0.3, 0.2), (0.5, 0.7) \rangle$ . Then  $M$  is a NS-CS but not a  $N_{\alpha g}$ -CS as  $M \subseteq A, B$  and  $N_{\alpha}Cl(M) = M \cup C(A) = C(A) \not\subseteq M$ .

**Remark 3.12:** Every NP-CS and every  $N_{\alpha g}$ -CS in a NTS  $X$  are independent to each other in general.

**Example 3.13:** In Example 3.11,  $M$  is a NP-CS but not a  $N_{\alpha g}$ -CS as seen in the respective example.

**Example 3.14:** Let  $X = \{a, b\}$  and  $\tau = \{0_N, A, B, 1_N\}$ , where  $A = \langle x, (0.5, 0.4), (0.3, 0.2), (0.5, 0.6) \rangle$  and  $B = \langle x, (0.4, 0.3), (0.3, 0.1), (0.6, 0.7) \rangle$ . Then  $\tau$  is a NT. Let  $M = \langle x, (0.5, 0.5), (0.2, 0.1), (0.4, 0.4) \rangle$ . Then  $M$  is a  $N_{\alpha g}$ -CS but not a NP-CS as  $NCl(NInt(M)) = C(A) \not\subseteq M$ .

**Proposition 3.15:** Every  $N_{\alpha}$ -CS  $A$  is a  $N_{\alpha g}$ -CS in  $X$  but not conversely in general.

**Proof:** Let  $A \subseteq U$ , where  $U$  is a NOS in  $X$ . Then  $N_{\alpha}Cl(A) = A \subseteq NCl(NInt(NCl(A))) \subseteq A \subseteq A = A \subseteq U$ , by hypothesis. Hence  $A$  is a  $N_{\alpha g}$ -CS in  $X$ .

**Example 3.16:** In Example 3.6,  $M$  is a  $N_{\alpha g}$ -CS in  $X$  but not a  $N_{\alpha}$ -CS as  $NCl(NInt(NCl(M))) = C(B) \not\subseteq M$ .

**Proposition 3.17:** Every NOS,  $N_{\alpha}$ -OS are  $N_{\alpha g}$ OS but not conversely in general.

**Proof:** Obvious.

**Example 3.18:** In Example 3.6,  $C(M)$  is a  $N_{\alpha g}$ OS but not a NOS,  $N_{\alpha}$ -OS in  $X$ .

**Remark 3.19:** Both NS-OS and NP-OS are independent to  $N_{\alpha g}$ OS in  $X$  in general.

**Example 3.20:** The above Remark can be proved easily from the Examples 3.10, 3.11 and 3.13, 3.14 respectively.

**Proposition 3.21:** The union of any two  $N_{\alpha g}$ CSs is a  $N_{\alpha g}$ CS in a NTS  $X$ .

**Proof:** Let  $A$  and  $B$  be any two  $N_{\alpha g}$ CSs in a NTS  $X$ . Let  $A \subseteq B \subseteq U$  where  $U$  is a NOS in  $X$ . Then  $A \subseteq U$  and  $B \subseteq U$ . Now  $N_{\alpha}Cl(A \cup B) = (A \cup B) \subseteq NCl(NInt(NCl(A \cup B))) \subseteq (A \cup B) \subseteq NCl(NCl(A \cup B)) \subseteq (A \cup B) \subseteq NCl(A \cup B) \subseteq NCl(A \cup B) = NCl(A) \cup NCl(B) \subseteq U \subseteq U = U$ , by hypothesis. Hence  $A \cup B$  is a  $N_{\alpha g}$ CS in  $X$ .

**Remark 3.22:** The intersection of any two  $N_{\alpha g}$ CSs need not be a  $N_{\alpha g}$ CS in a NTS  $X$ .

**Example 3.23:** Let  $X = \{a, b\}$  and  $\tau = \{0_N, A, B, 1_N\}$  where  $A = \langle x, (0.5, 0.4), (0.3, 0.2), (0.5, 0.6) \rangle$  and  $B = \langle x, (0.8, 0.7), (0.3, 0.2), (0.2, 0.3) \rangle$ . Then  $\tau$  is a NT. Let  $M = \langle x, (0.6, 0.9), (0.3, 0.2), (0.4, 0.1) \rangle$  and  $N = \langle x, (0.9, 0.7), (0.3, 0.2), (0.1, 0.3) \rangle$ . Then  $M$  and  $N$  are  $N_{\alpha g}$ CSs in  $X$  but  $M \cap N = \langle x, (0.6, 0.7), (0.3, 0.2), (0.4, 0.3) \rangle$  is not a  $N_{\alpha g}$ CS as  $M \cap N \subseteq B$  and  $N_{\alpha}Cl(M \cap N) = 1_N \not\subseteq A$ .

**Proposition 3.24:** Let  $(X, \tau)$  be a NTS. Then for every  $A \in N_{\alpha g}C(X)$  and for every  $B \in NS(X)$ ,  $A \subseteq B \subseteq N_{\alpha}Cl(A)$  implies  $B \in N_{\alpha g}C(X)$ .

**Proof:** Let  $B \subseteq U$  and  $U$  be a NOS in  $(X, \tau)$ . Then since  $A \subseteq B$ ,  $A \subseteq U$ . By hypothesis,  $B \subseteq N_{\alpha}Cl(A)$ . Therefore  $N_{\alpha}Cl(B) \subseteq N_{\alpha}Cl(N_{\alpha}Cl(A)) = N_{\alpha}Cl(A) \subseteq U$ , since  $A$  is an  $N_{\alpha g}$ CS in  $(X, \tau)$ . Hence  $B \in N_{\alpha g}C(X)$ .

**Proposition 3.25:** If  $A$  is a NOS and a  $N_{\alpha g}$ CS in  $(X, \tau)$ , then  $A$  is a  $N_{\alpha}$ -CS in  $(X, \tau)$ .

**Proof:** Since  $A \subseteq A$  and  $A$  is a NOS in  $(X, \tau)$ , by hypothesis,  $N_{\alpha}Cl(A) \subseteq A$ . But  $A \subseteq N_{\alpha}Cl(A)$ . Therefore  $N_{\alpha}Cl(A) = A$ . Hence  $A$  is a  $N_{\alpha}$ -CS in  $(X, \tau)$ .

**Proposition 3.26:** Let  $(X, \tau)$  be a NTS. Then every NS in  $(X, \tau)$  is a  $N_{\alpha g}$ CS in  $(X, \tau)$  if and only if  $N_{\alpha}-O(X) = N_{\alpha}-C(X)$ .

**Proof: Necessity:** Suppose that every NS in  $(X, \tau)$  is a  $N_{\alpha g}$ CS in  $(X, \tau)$ . Let  $U \in NO(X)$ . Then  $U \in N_{\alpha}-O(X)$  and by hypothesis,  $N_{\alpha}Cl(U) \subseteq U \subseteq N_{\alpha}Cl(U)$ . This implies  $N_{\alpha}Cl(U) = U$ . Therefore  $U \in N_{\alpha}-C(X)$ . Hence  $N_{\alpha}-O(X) \subseteq N_{\alpha}-C(X)$ . Let  $A \in N_{\alpha}-C(X)$ . Then  $C(A) \in N_{\alpha}-O(X) \subseteq N_{\alpha}-C(X)$ . That is  $C(A) \in N_{\alpha}-C(X)$ . Therefore  $A \in N_{\alpha}-O(X)$ . Hence  $N_{\alpha}-C(X) \subseteq N_{\alpha}-O(X)$ . Thus  $N_{\alpha}-O(X) = N_{\alpha}-C(X)$ .

**Sufficiency:** Suppose that  $N_{\alpha}-O(X) = N_{\alpha}-C(X)$ . Let  $A \subseteq U$  and  $U$  be a NOS in  $(X, \tau)$ . Then

$U \in N_{\square}\text{-O}(X)$  and  $N_{\square}\text{Cl}(A) \subseteq N_{\square}\text{Cl}(U) = U$ , since  $U \in N_{\square}\text{-C}(X)$ , by hypothesis. Therefore  $A$  is an  $N_{\square}\text{gCS}$  in  $X$ .

**Proposition 3.27:** If  $A$  is a NOS and a  $N_{\square}\text{gCS}$  in  $(X, \tau)$ , then  $A$  is a NROS in  $(X, \tau)$ .

**Proof:** Let  $A$  be a NOS and a  $N_{\square}\text{gCS}$  in  $(X, \tau)$ . Then  $A$  is a  $N_{\square}\text{-CS}$  in  $X$ . Now  $N\text{Int}(N\text{Cl}(A)) \square N\text{Cl}(N\text{Int}(N\text{Cl}(A))) \square A$ . Since  $A$  is a NOS,  $A = N\text{Int}(A) \square N\text{Int}(N\text{Cl}(A))$ . Hence  $N\text{Int}(N\text{Cl}(A)) = A$  and  $A$  is a NROS in  $X$ .

**Definition 3.28:** A NS  $A$  in  $(X, \tau)$  is a neutrosophic Q-set (NQ-S) in  $X$  if  $N\text{Int}(N\text{Cl}(A)) = N\text{Cl}(N\text{Int}(A))$ .

**Proposition 3.29:** For a NOS  $A$  in  $(X, \tau)$ , the following conditions are equivalent:

- (i)  $A$  is a NCS in  $(X, \tau)$ ,
- (ii)  $A$  is a  $N_{\square}\text{gCS}$  and a NQ-S in  $(X, \tau)$ .

**Proof:** (i)  $\Rightarrow$  (ii) Since  $A$  is a NCS, it is a  $N_{\square}\text{gCS}$  in  $(X, \tau)$ . Now  $N\text{Int}(N\text{Cl}(A)) = N\text{Int}(A) = A = N\text{Cl}(A) = N\text{Cl}(N\text{Int}(A))$ , by hypothesis. Hence  $A$  is a NQ-S in  $(X, \tau)$ .

(ii)  $\Rightarrow$  (i) Since  $A$  is a NOS and a  $N_{\square}\text{gCS}$  in  $(X, \tau)$ , by Theorem 3.27,  $A$  is a NROS in  $(X, \tau)$ . Therefore  $A = N\text{Int}(N\text{Cl}(A)) = N\text{Cl}(N\text{Int}(A)) = N\text{Cl}(A)$ , by hypothesis. Hence  $A$  is a NCS in  $(X, \tau)$ .

**Proposition 3.30:** Let  $(X, \tau)$  be a NTS. Then for every  $A \in N_{\square}\text{gO}(X)$  and for every  $B \in \text{NS}(X)$ ,  $N_{\square}\text{Int}(A) \subseteq B \subseteq A$  implies  $B \in N_{\square}\text{gO}(X)$ .

**Proof:** Let  $A$  be any  $N_{\square}\text{gO}$  of  $X$  and  $B$  be any NS of  $X$ . By hypothesis  $N_{\square}\text{Int}(A) \subseteq B \subseteq A$ . Then  $C(A)$  is a  $N_{\square}\text{gCS}$  in  $X$  and  $C(A) \subseteq C(B) \subseteq N_{\square}\text{Cl}(C(A))$ . By Theorem 3.24,  $C(B)$  is a  $N_{\square}\text{gCS}$  in  $(X, \tau)$ . Therefore  $B$  is a  $N_{\square}\text{gO}$  in  $(X, \tau)$ . Hence  $B \in N_{\square}\text{gO}(X)$ .

**Proposition 3.31:** Let  $(X, \tau)$  be a NTS. Then for every  $A \in \text{NS}(X)$  and for every  $B \in \text{NS-O}(X)$ ,  $B \subseteq$

$A \subseteq N\text{Int}(N\text{Cl}(B))$  implies  $A \in N_{\square}\text{gO}(X)$ .

**Proof:** Let  $B$  be a NS-OS in  $(X, \tau)$ . Then  $B \subseteq N\text{Cl}(N\text{Int}(B))$ . By hypothesis,  $A \subseteq N\text{Int}(N\text{Cl}(B)) \subseteq N\text{Int}(N\text{Cl}(N\text{Cl}(N\text{Int}(B)))) \subseteq N\text{Int}(N\text{Cl}(N\text{Int}(B))) \subseteq N\text{Int}(N\text{Cl}(N\text{Int}(A)))$ . Therefore  $A$  is a  $N_{\square}\text{-OS}$  and by Proposition 3.17,  $A$  is a  $N_{\square}\text{gO}$  in  $(X, \tau)$ . Hence  $A \in N_{\square}\text{gO}(X)$ .

**Proposition 3.32:** A NS  $A$  of a NTS  $(X, \tau)$  is a  $N_{\square}\text{gO}$  in  $(X, \tau)$  if and only if  $F \subseteq N_{\square}\text{Int}(A)$  whenever  $F$  is a NCS in  $(X, \tau)$  and  $F \subseteq A$ .

**Proof: Necessity:** Suppose  $A$  is a  $N_{\square}\text{gO}$  in  $(X, \tau)$ . Let  $F$  be a NCS in  $(X, \tau)$  such that  $F \subseteq A$ . Then  $C(F)$  is a NOS and  $C(A) \subseteq C(F)$ . By hypothesis  $C(A)$  is a  $N_{\square}\text{gCS}$  in  $(X, \tau)$ , we have  $N_{\square}\text{Cl}(C(A)) \subseteq C(F)$ . Therefore  $F \subseteq N_{\square}\text{Int}(A)$ .

**Sufficiency:** Let  $U$  be a NOS in  $(X, \tau)$  such that  $C(A) \subseteq U$ . By hypothesis,  $C(U) \subseteq N_{\square}\text{Int}(A)$ . Therefore  $N_{\square}\text{Cl}(C(A)) \subseteq U$  and  $C(A)$  is an  $N_{\square}\text{gCS}$  in  $(X, \tau)$ . Hence  $A$  is a  $N_{\square}\text{gO}$  in  $(X, \tau)$ .

**References:**

1. Atanassov, K., Intuitionistic fuzzy sets, Fuzzy Sets and Systems, 1986, 87- 96.
2. Coker, D., An introduction to intuitionistic fuzzy topological spaces, Fuzzy sets and systems, 88, 1997, 81-89.
3. Floretin Smarandache, Neutrosophic set:- A generalization of Intuitionistic fuzzy set, Jour. of Defense Resources Management, 2010, 107-116.
4. Ishwarya, P and Bageerathi, K., On Neutrosophic semi-open sets in Neutrosophic topological spaces, International Jour. of Math. Trends and Tech. 2016, 214-223.
5. Qays Hatem Imran, Smaandache, F., et. al., On Neutrosophic semi alpha open sets, Neutrosophic sets and systems, 2017, 37-42.
6. Salama, A. A., and Alblowi, S. A., Neutrosophic set and Neutrosophic topological spaces, IOSR Jour. of Mathematics, 2012, 31-35.